

THE CATIONS KrF^+ , XeF_3^+ , XeOF_3^+ AND XeOF_5^+
AND OXIDIZING PROPERTIES OF KrF^+

D. E. McKee, C. J. Adams, A. Zalkin, and Neil Bartlett

November 1972

RECEIVED
LAWRENCE
BERNARD LABORATORY


NOV 15 1972

LIBRARY AND
DOCUMENTS SECTION

Prepared for the U. S. Atomic Energy
Commission under Contract W-7405-ENG-48

For Reference

Not to be taken from this room



LBL-1163
c. |

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

The Cations KrF^+ , XeF_3^+ , XeOF_3^+ and XeOF_5^+ and Oxidizing Properties of KrF^+

By D. E. McKee, C. J. Adams, A. Zalkin, and Neil Bartlett

Inorganic Materials Research Division, Lawrence Berkeley Laboratory and Department of Chemistry; University of California, Berkeley, California 94720 U.S.A.

Summary - The XeOF_5^+ ion (prepared by exploiting KrF^+ as an oxidizer) is related to, but distinguishable from XeOF_3^+ ; the latter is in turn related to XeF_3^+ , the geometry of which has been established by X-ray crystallography.

In earlier communications¹ we showed that the adduct² $\text{KrF}_2 \cdot 2\text{SbF}_5$ was the salt $\text{KrF}^+\text{Sb}_2\text{F}_{11}^-$. The stretching frequency $\nu(\text{Kr-F}^+) = 626 \text{ cm}^{-1}$ is in excellent agreement with Schaefer's theoretical prediction³ of 620 cm^{-1} .

In accord with the anticipated⁴ high value of the electron affinity of KrF^+ , the cation has proved to be an extraordinary oxidizer. We have previously shown¹ that it oxidizes IF_5 to IF_6^+ and have now carried out the oxidation of XeOF_4 to XeOF_5^+ ($\text{XeOF}_4 + \text{KrF}^+ \rightarrow \text{XeOF}_5^+ + \text{Kr}$). Since the syntheses, which involve addition of XeOF_4 to $\text{KrF}^+\text{Sb}_2\text{F}_{11}^-$ (~ -10), have been executed in quartz apparatus, there has always been some contamination from O_2^+ salts, but the best preparations have yielded no XeOF_3^+ salts. Raman data for XeOF_5^+ (compared with data for IOF_5 and its Xe relatives in Figure 1) show a pattern of lines consistent with an IOF_5 -like species, but the Xe-F and Xe-O stretching frequencies are lower than for IOF_5 . As may be seen from Figure 1, this is akin to the situation in XeF_5^+ salts, where $\nu_{\text{symm}}(\text{Xe-F})$ tends to be somewhat lower than $\nu_{\text{symm}}(\text{I-F})$ in IF_5 .

As part of our study^{5,6,7} of the fluoride ion donor abilities of the xenon fluorides and oxyfluorides, we have also investigated the systems $\text{XeOF}_4/\text{SbF}_5$ and $\text{XeF}_4/\text{SbF}_5$: moreover, unambiguous identification of the XeOF_5^+ ion demanded a full characterization of the former system. A combination of X-ray crystallographic and Raman spectroscopic data has established that XeF_3^+ is the only cation present in the $\text{XeF}_4/\text{SbF}_5$ system; Raman spectroscopic evidence indicates that the XeOF_3^+ cation is the only one present in the $\text{XeOF}_4/\text{SbF}_5$ system. Since the onset of our studies Gillespie and his coworkers have given vibrational and ^{19}F nmr spectroscopic evidence for both XeF_3^+ (Ref. 8) and XeOF_3^+ (Ref. 9).

Our studies have shown that in the $\text{XeF}_4/\text{SbF}_5$ system there are two compounds, $\text{XeF}_3^+\text{SbF}_6^-$ and $\text{XeF}_3^+\text{Sb}_2\text{F}_{11}^-$: efforts to make $\text{Xe}_2\text{F}_7^+\text{SbF}_6^-$ have failed. Both salts are pale yellow-green solids. $\text{XeF}_3^+\text{SbF}_6^-$ (m.p. $109-113^\circ$) is dimorphic, with a transition temperature of $\sim 90^\circ$; the low temperature form is monoclinic with $a = 5.50$, $b = 15.50$, $c = 8.95$ ($a \pm 0.01 \text{ \AA}$), $\beta = 102.9 \pm 0.3^\circ$, $V = 743.3 \text{ \AA}^3$, $z = 4$, $D_c = 3.81 \text{ g cm}^{-3}$. $\text{XeF}_3^+\text{Sb}_2\text{F}_{11}^-$ (m.p. $81-83^\circ$) is triclinic with $a = 8.237(5)$, $b = 9.984(20)$, $c = 8.004(5)$, $\alpha = 72.54(5)$, $\beta = 112.59(7)$, $\gamma = 117.05(21)^\circ$, $V = 534.9 \text{ \AA}^3$, $z = 2$, $D_c = 3.98 \text{ g cm}^{-3}$. The structure of the latter has been successfully refined in space group PT using three-dimensional graphite-monochromatized MoK_α X-ray data. With anisotropic temperature factors for all atoms, a final conventional R factor of 0.035 for 1823 independent reflections for which $I \geq 3\sigma(I)$ has been obtained. The crystal structure is built up from the structural units, $\text{XeF}_3^+\text{Sb}_2\text{F}_{11}^-$, shown in Figure 2. The I shaped cation is planar and lies in the same plane as a fourth fluorine atom, which makes a close contact of 2.50 \AA

to the xenon atom. This interaction of the cation and the anion is consistent with a distorted trigonal bipyramidal configuration of two axial F ligands, one equatorial F ligand and two sterically active, equatorial, non-bonding valence-electron pairs about the Xe atom. Such a cation should have a maximum polarizing effect normal to the triangular faces containing the two non-bonding pairs. As in the electronically related molecules ClF_3 (Ref. 10) and BrF_3 (Ref. 11) the axial bonds in XeF_3^+ (1.88 and 1.89 Å) are longer than the equatorial (1.83 Å). This is consistent with designation of the latter as an electron-pair bond and the former as three-center four-electron bonds^{4,12}.

In the $\text{XeOF}_4/\text{SbF}_5$ system, Selig had previously established¹³ the compound $\text{XeOF}_4 \cdot 2\text{SbF}_5$, but structural information was lacking until the recent report by Gillespie and his coworkers⁹. In our studies, two compounds have been isolated (1:1, m.p. 104-105° and 1:2, m.p. 61-66°). The XeOF_4 , in the SbF_5 complexes, is certainly no longer molecular, as in the 1:1 $\text{XeF}_2 \cdot \text{XeOF}_4$ molecular adduct¹², and the marked increase in the Xe-F stretching frequency clearly evinces cation formation. Furthermore, comparisons of the Raman spectra of the SbF_5 complexes, given in Figure 1, indicate the salt formulations $\text{XeOF}_3^+ \text{SbF}_6^-$ and $\text{XeOF}_3^+ \text{Sb}_2\text{F}_{11}^-$. The similarities of the XeOF_3^+ and XeF_3^+ spectra suggest a close structural relationship. It is therefore probable that the XeOF_3^+ geometry will be very like that of XeF_3^+ to which an oxygen atom has been added at a Xe electron-pair site (equatorial).

The similarity of the Xe-O stretching frequencies suggests that the Xe-O bonds in XeOF_3^+ and XeOF_4 must be nearly the same. Also, that the axial stretching frequencies of XeOF_3^+ lie somewhat higher than

those of XeF_3^+ indicates that the Xe-F axial bonds will be slightly shorter in the former than in the latter.


This work was supported by the U.S. Atomic Energy Commission and CJA is grateful to the Commonwealth fund for a Harkness Fellowship.

References

1. D. E. McKee, R. Mews, and N. Bartlett, Abstracts 162nd American Chemical Society National Meeting, September 14-16, 1971, FLUO 2; and N. Bartlett, D. Gibler, D. McKee, R. Mews and M. Wechsberg, Abstracts of The Sixth International Symposium on Fluorine Chemistry, July 18-23, 1972, C24.
2. H. Selig and R. D. Peacock, J. Amer. Chem. Soc. 1964, 86, 3895.
3. B. Liu, H. F. Schaefer III, J. Chem. Phys. 1971, 55, 2369.
4. N. Bartlett, Endeavour, 1972, XXXI, 107.
5. F. O. Sladky, P. A. Bulliner, N. Bartlett, B. Y. DeBoer, and A. Zalkin, Chem. Comm., 1968, 1048; and F. O. Sladky, P. A. Bulliner and N. Bartlett, J. Chem. Soc. (A), 1969, 2180.
6. N. Bartlett and F. O. Sladky, J. Amer. Chem. Soc., 1968, 90, 5316.
7. N. Bartlett, M. Gennis, D. D. Gibler, B. K. Morrell, and A. Zalkin. (LBL Report No. 850) submitted for publication in Inorg. Chem.
8. R. J. Gillespie, B. Landa, and G. J. Schrobilgen, Chem. Comm., 1971, 1543.
9. R. J. Gillespie, B. Landa, and G. J. Schrobilgen, Chem. Comm., 1972, 607.

10. D. F. Smith, J. Chem. Phys., 1953, 21, 609.
11. D. W. Magnuson, J. Chem. Phys., 1957, 27, 223.
12. N. Bartlett and M. Wechsberg, Z. anorg. Chem., 1971, 385, 5.
13. H. Selig, Inorg. Chem., 1966, 5, 183.
14. L. E. Alexander and I. R. Beattie, J. Chem. Soc. (A), 1971, 3091.
15. D. F. Smith and G. M. Begun, J. Chem. Phys., 1965, 43, 2001.
16. C. J. Adams and A. J. Downs, Spectrochim. Acta, 1972, 28A, 1841.

Figure 1: Raman Spectra of Xenon-Cation Species and Iodine Relatives



Footnotes for Figure 1.

* Raman line assigned to anion.

(a) Ref. 14.

(b) C. J. Adams and N. Bartlett, to be published. The fundamental frequencies of XeF_5^+ are sensitive functions of the counterion and phase, although solid $\text{XeF}_5^+\text{BF}_4^-$ is a typical example. Common values for the stretching frequencies are: $\nu_1(a_1)$ 650-680 cm^{-1} ; $\nu_2(a_1)$ 585-630 cm^{-1} ; $\nu_4(b_1)$ 600-640 cm^{-1} ; $\nu_7(e)$ 640-670 cm^{-1} .


(c) Ref. 15.

(d) This work. Lines attributable to dioxygenyl fluoroantimonates have been deleted.

(e) This work. Assignments for XeOF_3^+ : 944 cm^{-1} , $\nu(\text{Xe-O})$; 649, ν_{asym} (ax. XeF_2); 638, $\nu(\text{eq. Xe-F})$; 601 ν_{symm} (ax. XeF_2); 358, 333, $\delta(\text{FXeF})$, $\delta'(\text{OXeF})$. Our assignments differ from those of Gillespie et al. (Ref. 9) only in placing ν_{asym} (ax. XeF_2) higher than ν_{symm} (ax. XeF_2), as has been found for structurally related molecules, e.g. TeF_4 (Ref. 16).

(f) This work. Assignments for XeF_3^+ : 640 cm^{-1} , $\nu(\text{eq. Xe-F})$; 618 ν_{asym} (ax. XeF_2); 582 ν_{symm} (ax. XeF_2); 363 $\delta(\text{FXeF})$.

NOTE TO PRINTER: Figure 2 comprises the drawing (page 9) and the table (page 10).



XBL 729-6975

Figure 2: The $\text{XeF}_3^+\text{Sb}_2\text{F}_{11}^-$ structure unit.

<u>Distances (Å)</u>	<u>Angles (Deg.)</u>
Xe-Fe(2) 2.50(1)	F(9)-Xe-F(10) 81.73(30)
Xe-F(9) 1.83(1)	F(9)-Xe-F(11) 80.22(30)
Xe-F(10) 1.88(1)	F(10)-Xe-F(11) 161.90(40)
Xe-F(11) 1.89(1)	F(10)-Xe-F(2) 72.67(27)
	F(9)-Xe-F(2) 154.39(38)
Sb(1)-F(1) 2.01(1)	
Sb(1)-F(2) 1.90(1)	Xe-F(2)-Sb(1) 171.64(13)
Sb(1)-F(3) 1.85(1)	Sb(1)-F(1)-Sb(2) 155.37(15)
Sb(1)-F(5) 1.84(1)	
Sb(1)-F(8) 1.85(1)	
Sb(1)-F(12) 1.83(1)	

Figure 2: The $\text{XeF}_3^+\text{Sb}_2\text{F}_{11}^-$ Structure unit.

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720